

Urban heat and residential electricity consumption

Antunes De Azevedo, Juliana; Chapman, Lee; Muller, Catherine

DOI:

[10.1016/j.apgeog.2016.03.002](https://doi.org/10.1016/j.apgeog.2016.03.002)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Antunes De Azevedo, J, Chapman, L & Muller, C 2016, 'Urban heat and residential electricity consumption: a preliminary study', *Applied Geography*, vol. 70, pp. 59-67. <https://doi.org/10.1016/j.apgeog.2016.03.002>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Urban heat and residential electricity consumption: a preliminary study.

Abstract

The Urban Heat Island (UHI) is a well-documented phenomenon occurring in cities across the world resulting in city centres often being several degrees warmer than their surroundings. This local elevation in temperatures could potentially impact upon local energy consumption, with residents in the warmer central section of the city using more energy to cool their homes in summer and less energy to warm them in winter. This study uses a combination of Geographical Information System techniques and Remote Sensing data (MODIS LST and NDVI), as a preliminary investigation, to assess the spatial relationship between UHI, urban greenspace, household income and electricity consumption in Birmingham, UK. It provides simple and repeatable steps, based on freely available datasets, for urban planners, industry, human and physical geographers, and non-specialists to reproduce the analyses. The results show that, the present impact of the UHI is limited and instead highlights the dominance of household income over local climate in explaining consumption patterns across Birmingham.

Key words: Residential Electricity Consumption; Urban Greenspace; Income; Urban Heat Island.

1. Introduction

Energy demand in urban areas is an important facet of energy supply planning. In particular, increasing energy consumption by the residential sector is an issue that could endanger broader economic development since in itself it does not generate wealth and could limit the amount of energy available for other productive sectors (Pereira and Assis 2013). The electricity consumption by sectors in the UK can be observed in Figure 1, domestic consumption has maintained itself as the larger consuming sector almost throughout the whole period from 1965 to 2013.

[Figure 1 near here]

Household size, income, building design characteristics and local climatic conditions are all key factors in determining residential energy consumption (Santamouris et al. 2007). Generally, small households need less energy due to a reduced transfer area, but they also have lower occupancy, and therefore, fewer appliances when compared with larger households (Pérez-Lombard, Ortiz, and Pout 2008). Similarly, household income is an important factor, with a strong correlation evident between daily electricity consumption and earnings (Ghisi, Gosch, and Lamberts 2007). This pattern is evident spatially, where areas with higher average per capita income consume considerably more energy; a direct result of the relationship between energy consumption and the purchasing power of families (Pereira and Assis 2013).

With respect to climatic factors, the Urban Heat Island (UHI) is a potentially important localised phenomenon to take into account when assessing consumption in cities. The UHI is

described as the difference in temperature between an urban area and the surrounding rural area of the conurbation. It is mainly caused by anthropogenic changes to the environment with a range of factors contributing such as urban geometry, density / population of a conurbation, replacement of vegetation cover by construction material (e.g. asphalt and concrete), changing surface's albedo and emissivity thus reducing evapotranspiration and increased emissions of anthropogenic heat (Oke 1987). The overall result is that cities are generally warmer than their rural surroundings, reaching a maximum under "ideal" conditions (e.g., clear skies and light winds). For small cities, the effect can be minimal, for example, differences of 1°C in Ljutomer, Slovenia (Ivajnsič, Kaligarič, and Žiberna 2014) where as differences of more than 7 °C are not uncommon for large cities (e.g. Paris, France: (Lac et al. 2013). The subtle changes in temperature caused by UHI can impact on many aspects of everyday life, such as critical infrastructure (Chapman, Azevedo, and Prieto-Lopez 2013), health (Tomlinson et al. 2011) and energy consumption (Santamouris et al. 2001), with such impacts becoming exacerbated under heatwave events. It is hypothesised that the UHI should have a direct impact on energy consumption, particularly in the warmer core of the city (Taha et al. 1988, Kolokotroni et al. 2010, Hassid et al. 2000, Akbari, Pomerantz, and Taha 2001) where higher energy loads will be required for cooling in summer, and in winter consumption will reduce for heating. For example, in centrally located buildings in Athens (Greece), where the average UHI can exceed 10°C, cooling loads can double in summer, whereas winter period heating loads can decrease by 30% (Santamouris et al. 2001). Therefore, by not considering the UHI, energy consumption and peak power should be significantly underestimated (Hassid et al. 2000).

Green spaces are a widely adopted strategy to mitigate UHI intensity (Lambert-Habib et al. 2013) since they reduce urban temperatures thorough evapotranspiration and shadowing. In

modelling experiments carried out for Manchester, UK, it was found that a 5% increase in mature deciduous trees can reduce average hourly surface temperatures by 1°C during summer (Skelhorn, Lindley, and Levermore 2014). For example, the highest cooling loads in Athens are seen in the western area of the city where there is limited greenspace (Santamouris et al. 2001). In Manchester, it is proposed that if all vegetation was replaced with asphalt, then air temperature would increase by up to 3.2°C at midday (Skelhorn, Lindley, and Levermore 2014). Similarly, it was found in the USA that for an increase of 25% of tree cover in urban areas can result in a 40% annual residential cooling energy savings in Sacramento and 25% in Phoenix and Lake Charles (Huang et al. 1987).

The UHI can be largely subdivided into three different types: the surface UHI, the canopy UHI (i.e. 2m) and the boundary layer UHI (Azevedo, Chapman, and Muller 2016). Air temperature measurements are used to quantify both the the canopy and boundary UHI, whereas land surface temperature (LST) is used for the surface UHI. Traditional ways in which canopy UHI are measured include station pairs (e.g. Wilby 2003) or the use of transects (e.g. Smith et al. 2011). However, these usually have limited spatial (Smith et al. 2011, Muller et al. 2013) and temporal resolution, and therefore there has been an ongoing challenge to quantify the intensity and spatial extent of the canopy UHI.

Due to the wide spatial coverage and availability of data, thermal remote sensing is one of the most popular techniques used for the evaluation of UHI (Roth, Oke, and Emery 1989, Tomlinson, Chapman, Thornes, and Baker 2012, Smith et al. 2011, Yuan and Bauer 2007, Weng, Lu, and Schubring 2004, Schwarz, Lautenbach, and Seppelt 2011, Keramitsoglou et al. 2011, Dousset 1989, Dousset, Laaidi, and Zeghnoun 2011, Azevedo, Chapman, and Muller 2016). The main advantage is that remote sensing provides a consistent, repeatable

methodology for the end-user (Tomlinson et al. 2011). However, thermal remote sensing observes LST which restricts studies to just the surface UHI. Although LST plays a major role in urban climatological processes as surface temperature modulates the air temperature of the urban canopy layer, and therefore influences the internal climate of buildings and general thermal comfort (Voogt and Oke 2003), it can only provide an indication of air temperatures and therefore the canopy UHI. Furthermore, remote sensing isn't ideal to evaluate the UHI in small cities, since the spatial resolution of the sensor can often be too coarse (Ivajnsič, Kaligarič, and Žiberna 2014).

Vegetation abundance is an influential factor controlling UHI (Weng, Lu, and Schubring 2004) and the Normalized Differenced Vegetation Index (NDVI) is often used to approximate vegetation abundance. The connection between NDVI and LST has been well established in studies, and a negative relationship between NDVI and LST has been shown and proven to be seasonally variable (e.g. Yuan and Bauer 2007). Other studies which have included energy consumption data in the analysis (Akbari, Pomerantz, and Taha 2001, Huang et al. 1987), but no study has yet investigated all these factors along with income and socioeconomic data at the same temporal and spatial resolution (e.g. Pereira and Assis 2013, Santamouris et al. 2007). Hence, this article aims to combine energy and income data with LST and NDVI data to assess the relationship between income, UHI¹, vegetation and residential electricity consumption in Birmingham, UK. It also focuses on simple and repeatable steps, based on freely available datasets. The results could be used to inform current residential electricity consumption modelling due to the UHI effect.

2. Study area

¹ Since LST data is being analysed, only surface UHI is being addressed, however the general term UHI will be used.

Birmingham is the second largest urban area in the UK with an estimated population of over 1 million people (Birmingham City Council 2014). It is a post-industrial city with a distinct range of land use zones (e.g. the central business district, eastern industrial areas with the majority of residential areas straddling this belt of commerce and industry to the north and south). Some large parks can be found closer to the wealthier neighbourhoods (Figure 2a).

[Figure 2 near here]

Tomlinson, Chapman, Thornes, and Baker (2012) used night-time MODIS imagery for the summer of 2003-2009 and identified that during periods of high atmospheric stability, the surface UHI magnitude in Birmingham can reach up to 5-7°C. The cooling effect of large areas of greenspace in Birmingham was evident, particularly in Sutton Park, Woodgate Valley and the Lickey Hills (mentioned locations in Figure 2b), with a significant temperature gradient extending northwards from the city centre to Sutton Park (~ distance of 10 km) where temperatures can be 7-8°C cooler than the urban core under heatwave conditions (Tomlinson et al. 2013).

3. Methodology, Datasets & Analysis

3.1. Electricity consumption and income data

Ordinary residential electricity consumption data and income model based estimates are available from the UK Department of Energy and Climate Change (DECC) and the UK Office for National Statistics (ONS) respectively. Both datasets are aggregated into Super

Output Areas (SOAs), a standard unit used in the UK to report areal statistics (although any areal statistic unit is viable to reproduce the work elsewhere). SOAs don't have consistent physical size, but are instead based on established ranges of population and households for Census purposes (Table 1 – ONS 2011a). Income data is not available for the lower level (LSOA), hence middle level data (MSOA) is the universal unit considered for this study.

[Table 1 near here]

The fact that SOAs do not have a consistent physical size can raise questions regarding the stability of the estimates. Indeed, the spatial aggregation processing of geographical units have been extensively reviewed, and a number of different techniques are available to overcome bias (Jacobs-Crisioni, Rietveld, and Koomen 2014). For example, Bayes adjustment (Assunção et al. 2005) is a possible means to overcome the problems related to the demographic data, however the approach would not be applicable to the other data used in this study. Furthermore, the stability of the unit areas from one Census to the next is a known problem when using Census units (Fotheringham and Wong 1991). Despite these concerns, such units continue to be used in scientific studies and remain effective for spatial risk assessments being applicable to both the scale and preliminary focus of the research (Tomlinson et al. 2011, Pereira and Assis 2013).

Three types of electricity data are available from DECC (DECC 2013) recorded as total consumed over a year; Economy 7, Ordinary electricity consumption and Total electricity consumption. Economy 7 is a cheaper tariff (NB: this tariff is unique to the UK, other countries might or might not have similar alternatives) which offers the opportunity for users to concentrate their usage during a 7 hour period at night (for example, the charging of night

storage heaters) where as ordinary consumption is the reminder of other tariffs. Total electricity consumption is simply the combination of the two. This study considers only ordinary energy consumption data as Economy 7 has a tendency to be used independently of weather as it lacks the ‘controllability’ of other tariffs – i.e. a ‘set point’ where users turn on heating and cooling systems. For the analysis, MSOA consumption data for 2006 was used normalized by the number of households. Firstly, a simple normalization through division was performed indicating the average consumption (the total ordinary consumption by MSOA) by household (number of households by MSOA). Secondly, MSOA consumption by household was normalized by the household income.

With respect to income data, the ONS income estimate model has a 95% confidence level and estimates households average weekly income. Model based income estimates per MSOA for 2007/2008 were used (the closest to 2006 - other releases are 2001/2002, 2004/2005, and 2011/2012) (ONS 2011b).

3.2. LST data

Satellite data, also from 2006, was aggregated to produce an annual summary. LSTs were analysed for both daytime and night-night for cloudless conditions to evaluate general UHI pattern for the year. Absolute temperatures values were used and considered to be more appropriate than residual temperatures for the analyses in this paper. Data was obtained from MODIS Aqua, with an overpass in the study area, ~01:30 and ~13:30 (UTM). The product used was MYD11A1 (V5) – MODIS/Aqua Land Surface Temperature and Emissivity Daily L3 Global 1km Grid SIN (LPDAAC 2015a). This product does present a compromise in spatial resolution (e.g. when compared to the 60m thermal band of Landsat 7 - i.e. after

February 25th, 2010, Landsat 7 thermal band is collected at 60m but resampled to 30m (USGS 2010)), but the vastly improved temporal resolution of Aqua greatly increases image availability for the study. This is an important factor for studies in the UK, where cloud cover is a frequent and prohibitive problem when using Landsat, which overpasses the study area just once every 16 days.

The MODIS LST product uses a split window algorithm to correct for atmospheric effects (LPDAAC 2015a) and surface emissivity (Tomlinson, Chapman, Thornes, and Baker 2012) and has been used in Birmingham in previous studies (Tomlinson, Chapman, Thornes, Baker, et al. 2012, Tomlinson, Chapman, Thornes, and Baker 2012, Azevedo, Chapman, and Muller 2016). The MODIS Reprojection Tool (MRT) (LPDAAC 2014) was used to convert images to GeoTIFF format at UTM, and subsequently converted to British National Grid (BNG) in ArcGIS (MODIS products are released at Sinusoidal Projection). For the night-time analysis, 45 cloud free images were available and for daytime, 27 images were retained. In both cases the largest amount of images available were during summer and autumn, because of the more stable weather conditions on those seasons with decreasing cloud cover. Data averaging and quality control was then conducted in ArcGIS, where the final 100% cloud free images were selected; before being converted from Kelvin to Celsius, and clipped to the study area. The result was one averaged image for daytime LST (Figure 3a) and one for night-time LST (Figure 3b).

3.3. NDVI dataset

The NDVI dataset was also obtained from Aqua MODIS products, MYD13Q1 (V5) – MODIS/Aqua Vegetation Indices 16-Day L3 Global 250m Grid SIN (LPDAAC 2015b),

available in Sinusoidal Projection, every 16 days at 250 m resolution. The product is the difference between pigment absorption features in bands 1 (red reflectance) and 2 (near infrared). It is atmosphere-corrected and quality controlled, based on a 16 day composite (LPDAAC 2015b). Two vegetation indices are available for each product NDVI and EVI (Enhanced Vegetation Index). EVI was not used in this study since it is more applicable to monitor changes in canopy structure and leaf area, whereas NDVI is used to verify vegetation density and is the index most frequently used by urban climate studies (Weng, Lu, and Schubring 2004, Yuan and Bauer 2007). NDVI ranges from -1 to 1, being positive values increasing amount of vegetation in a pixel (Yuan and Bauer 2007), while 0 and negative values indicate rock, asphalt, clouds, snow, ice and water.

As per the LST product, the data was downloaded and converted in MODIS MRT to GeoTIFF format at UTM, and subsequently to British National Grid (BNG) in ArcGIS. All NDVI images available for 2006 were used, resulting in 23 images for the study period (one every 16 days). ArcGIS was then used to apply a scale factor (as indicated in reference material - (LPDAAC 2015b)) to adjust the range from -1 to 1. Finally, the 23 images were averaged into a single image for the year and clipped to the Birmingham area (Figure 3c).

[Figure 3 near here]

3.4. Data aggregation and analysis

As income and residential electricity consumption data is available by MSOA, for analysis purposes, there was a need to average and aggregate LST and NDVI into MSOAs (Figure 4). The processed LST and NDVI raster images were simply summed and then averaged by the

number of images used before being converted into a point dataset. All points located within each MSOA were then averaged, resulting in a unique LST or NDVI value by MSOA. Direct correlations between the variables was then calculated by using Pearson correlation coefficients (Table 2). Direct Pearson correlation was also carried between the MSOA consumption by household normalized by the income data and the aggregated LST and NDVI by MSOAs (Table 3). *P-values* lower than 0.01 were found for all correlations; considering a standard $\alpha = 0.05$ cut off, all analyses are significant at a 95% confidence interval. Scatter plot diagrams with intercept, slope and R^2 are shown in Figure 5.

[Table 2 near here]

[Table 3 near here]

[Figure 4 near here]

[Figure 5 near here]

4. Discussion

All MODIS products used in this analyses were obtained for free online (USGS 2013) and are available since 2002 with worldwide coverage. The advantage of using the MODIS Aqua dataset is that is well-suited to non-specialists due to the fact that it is already atmospheric corrected with NDVI already calculated. Add to this, the fact that is free and available either twice a day for the LST, or in a 16 days composite for NDVI, it allows the user to determine the temporal scale of the study being carried, for yearly period, seasonally, monthly or daily.

Census data are usually available in most countries and free, which provides demographic investigation data and areal units. Other variables, sophisticated datasets, and areal units can be used for the analyses, depending on the scope and aim of the study and availability.

Although the UK electricity data has good spatial resolution, data is only available as an annual summary per MSOA and therefore doesn't allow for seasonal interpretation. Indeed, this can be seen as a problem, since the correlation between climate and electricity consumption has different patterns during summer and winter, and so does the UHI pattern and vegetation. However, for preliminary investigation focusing on freely available datasets at the same spatial and temporal resolution to provide results for spatial risk assessment it can still be used, and it is simple and repeatable.

As per Tomlinson et al. (2011), a clear UHI is evident in the averaged LST data with temperatures peaking in the city centre and significantly lower LST in the urban greenspace (Figures 3a and 3b). The range of LST evident during the day is higher than during the night, a consequence of differential solar heating of surfaces with different thermal properties during the daytime. After sunset surfaces start releasing energy absorbed during the day, cooling down. In the second case, air temperature is usually higher than LST.

The averaged NDVI distribution for Birmingham (Figure 3c) was also as expected ranging from 0.2 in the city centre to 0.7 in the larger urban greenspaces. As demonstrated in previous studies (Weng, Lu, and Schubring 2004; Yuan and Bauer 2007) a strong negative correlation between LST and NDVI exists, with the strongest relationship evident during the daytime ($r = -0.78$ compared to $r = -0.69$ at night). Furthermore, there is a strong positive correlation between income and NDVI ($r = 0.61$) and is explained by increased real estate

values surrounding parks and greenspace (Lambert-Habib et al. 2013). It is evident that wealthier families and individuals usually live in more vegetated areas (e.g. Sutton Coldfield in Birmingham); whereas lower income groups live in flats in cheaper areas (Santamouris et al. 2007), often close to the city centre (e.g. Ladywood in Birmingham). In Birmingham, it is not uncommon to find low-income groups living in areas where the UHI reaches its maximum, which when factored in with the poor housing stock found in such areas (i.e. less efficient construction and insulation), has implications for not only energy consumption but the general wellbeing and health of the population in these areas (Tomlinson, Chapman, Thornes, and Baker 2012). The same was found for Athens (Santamouris et al. 2007), however such statement should be analysed individually depending on the city studied, due to differences in culture, urban form and development of cities across the world.

The strongest relationship found with electricity consumption was with income ($r = 0.62$), highlighting that although low income groups have a greater need for heating (less well insulated housing stock) and air conditioning (increased exposure to UHI), the main driver for consumption is purchasing power (Table 2 and 4). It is primarily for this reason why direct correlations with LST are weaker than would be expected (daytime $r = -0.47$; night-time $r = -0.43$). It was hypothesised that higher temperatures in the city would result in increases electricity consumption by the use of air conditioners and fans, however that was not clearly identified in the first part of the analysis (Table 2).

[Table 4 near here]

As a second part of the analysis, the electricity consumption data was normalized by income to attempt to isolate the UHI influence in the electricity consumption, especially with daytime

UHI (Table 3). The correlation with the normalized consumption was higher for daytime UHI (daytime $r = 0.52$; night-time $r = 0.41$). A negative correlation with NDVI was observed ($r = -0.53$). In this case, all correlations marginally improved this analysis, but are still limited. Income may be an indirect factor of household size in this case, however such analyses are beyond the scope of this study.

Although the relationships were consistently significant, low correlations were obtained in some of the analyses (Figure 5). This can be attributed to other factors that may influence consumption, as well as the resolution and aggregation level of the variables. There is a clear need for other variables and data with higher spatial and temporal resolution to be taken into account in future and more detailed research. However, despite these limitations, it is evident that income is the most influential factor in electricity consumption, potentially an indirect factor of household size. The UHI appears to play a role, but these results are presently tempered and even with the presence of a strong UHI, high temperatures are still not an issue in Birmingham, therefore there is actually no significant need for cooling appliances at the moment.

5. Conclusions and Final Remarks

Despite electricity consumption data not being available at the desired temporal scale, it was possible to assess residential electricity consumption distribution and its correlation with income, NDVI and LST for yearly aggregated data, at a preliminary stage, based on simple and repeatable steps with freely available datasets. Large differences are evident in the distribution of urban heat and vegetation across Birmingham, but the results show that the dominant factor that influences residential electricity consumption at these scales is not

climate but income. Whether this is true at other scales is difficult to assess given the present spatial and temporal limitations of the available data. From this study, it would be easy to conclude that electricity consumption due to increasing temperatures does not seem like a current or urgent issue in temperate countries, however considering climate change scenarios, an increasing frequency of heatwaves and energy security concerns, overlooking behavioural changes of the millions of people who live in mid-latitude cities would be an oversight. In face of climate change scenarios in Birmingham, temperatures will increase (Azevedo, Chapman, and Muller 2015), exacerbating the UHI effect and impacts on electricity consumption. Also, the increasing number of people in urban areas will not only contribute to the exacerbation of the UHI effect but will also increase the number of people exposed to its potential risks (Smith et al. 2011), therefore, overlooking increasingly important climate drivers would be foolhardy.

Higher resolution data would certainly aid analysis, and with the advent of smart metering (i.e. the Internet of Things), consumption data will soon become available at seasonal and even daily scales, allowing better interpretation of LST, NDVI and income with consumption. The Internet of Things is now providing unparalleled opportunities for high resolution weather monitoring in our cities (e.g. crowdsourcing for weather and climate information:(Muller et al. 2015). Additionally, high resolution vegetation maps and land use maps can be freely extracted from Google Earth Imagery, and used to infer the impacts of different types of greening on temperature. Finally, the emerging Volunteered Geographic Information (Goodchild 2007, Arribas-Bel 2014), is available to the aid acquisition and validation of geographical information and variables to improve analyses (Foody et al. 2013, Basiouka and Potsiou 2012, Hawthorne et al. 2015, Spinsanti and Ostermann 2013). Hence, moving forward, there is tremendous potential for future research to understand electricity

consumption and urban climate at a high temporal and spatial resolution for spatial risk assessment, urban planning and energy industry, for current and future scenarios

Acknowledgments

The authors would like to thank the Brazilian agency CNPq (Brazilian National Council for Scientific and Technological Development – Research Grant: [236849/2012-3]) for their financial support of this research. The research would not have been possible without satellite data from NASA and U.S. Geological Survey, small area income estimates from the Office for National Statistics and energy estimates from the Department of Energy and Climate Change. A special thanks to Dr. Charlie Tomlinson who processed the original MODIS LST dataset.

References

- Akbari, H., M. Pomerantz, and H. Taha. 2001. "Cool surfaces and shade trees to reduce energy use and improve air quality in urban areas." *Solar Energy* no. 70 (3):295-310. doi: [http://dx.doi.org/10.1016/S0038-092X\(00\)00089-X](http://dx.doi.org/10.1016/S0038-092X(00)00089-X).
- Arribas-Bel, Daniel. 2014. "Accidental, open and everywhere: Emerging data sources for the understanding of cities." *Applied Geography* no. 49:45-53.
- Assunção, Renato M, Carl P Schmertmann, Joseph E Potter, and Suzana M Cavenaghi. 2005. "Empirical Bayes estimation of demographic schedules for small areas." *Demography* no. 42 (3):537-558.
- Azevedo, Juliana Antunes, Lee Chapman, and Catherine Muller. 2016. "Quantifying the Daytime and Night-Time Urban Heat Island in Birmingham, UK: A Comparison of Satellite Derived Land Surface Temperature and High Resolution Air Temperature Observations." *Remote Sensing* no. 8 (2):153.
- Azevedo, Juliana Antunes, Lee Chapman, and Catherine L. Muller. 2015. "Critique and suggested modifications of the degree days methodology to enable long-term electricity consumption assessments: a case study in Birmingham, UK." *Meteorological Applications* no. 22 (4):789-796. doi: 10.1002/met.1525.
- Basiouka, S, and C Potsiou. 2012. "VGI in Cadastre: a Greek experiment to investigate the potential of crowd sourcing techniques in Cadastral Mapping." *Survey Review* no. 44 (325):153-161.

- BCC, Birmingham City Council. *Population in Birmingham* 2014. Available from <http://www.birmingham.gov.uk/>.
- Chapman, Lee, Juliana Antunes Azevedo, and Tatiana Prieto-Lopez. 2013. "Urban heat & critical infrastructure networks: A viewpoint." *Urban Climate* no. 3:7-12. doi: <http://dx.doi.org/10.1016/j.uclim.2013.04.001>.
- DECC, Department of Energy and Climate Change. *Lower and Middle Super Output Areas electricity consumption* 2013. Available from <https://www.gov.uk/government/organisations/department-of-energy-climate-change>.
- Dousset, Bénédicte 1989. AVHRR-derived cloudiness and surface temperature patterns over the Los Angeles area and their relationship to land use. Paper read at Proceedings of IGARSS-89.
- Dousset, Bénédicte , Karine Laaidi, and Abdelkrim Zeghnoun. 2011. Surface temperature variability and mortality impact in the Paris region during the August 2003 heat wave. *Urban Climate News, Quarterly newsletter of International Association of Urban Climate* (42): 7-14
- Footy, Giles M, L See, Steffen Fritz, Marijn Van der Velde, Christoph Perger, Christian Schill, and Doreen S Boyd. 2013. "Assessing the accuracy of volunteered geographic information arising from multiple contributors to an internet based collaborative project." *Transactions in GIS* no. 17 (6):847-860.
- Fotheringham, A Stewart, and David WS Wong. 1991. "The modifiable areal unit problem in multivariate statistical analysis." *Environment and planning A* no. 23 (7):1025-1044.
- Ghisi, Eneidir, Samuel Gosch, and Roberto Lamberts. 2007. "Electricity end-uses in the residential sector of Brazil." *Energy Policy* no. 35 (8):4107-4120. doi: <http://dx.doi.org/10.1016/j.enpol.2007.02.020>.
- Goodchild, Michael F. 2007. "Citizens as sensors: the world of volunteered geography." *GeoJournal* no. 69 (4):211-221. doi: 10.1007/s10708-007-9111-y.
- Hassid, S., M. Santamouris, N. Papanikolaou, A. Linardi, N. Klitsikas, C. Georgakis, and D. N. Assimakopoulos. 2000. "The effect of the Athens heat island on air conditioning load." *Energy and Buildings* no. 32 (2):131-141. doi: [http://dx.doi.org/10.1016/S0378-7788\(99\)00045-6](http://dx.doi.org/10.1016/S0378-7788(99)00045-6).
- Hawthorne, T. L., V. Elmore, A. Strong, P. Bennett-Martin, J. Finnie, J. Parkman, T. Harris, J. Singh, L. Edwards, and J. Reed. 2015. "Mapping non-native invasive species and accessibility in an urban forest: A case study of participatory mapping and citizen science in Atlanta, Georgia." *Applied Geography* no. 56:187-198. doi: <http://dx.doi.org/10.1016/j.apgeog.2014.10.005>.
- Huang, Y. J., H. Akbari, H. Taha, and A. H. Rosenfeld. 1987. "The Potential of Vegetation in Reducing Summer Cooling Loads in Residential Buildings." *Journal of Climate and Applied Meteorology* no. 26 (9):1103-1116. doi: 10.1175/1520-0450(1987)026<1103:TPOVIR>2.0.CO;2.
- Ivajnsiĉ, Danijel, Mitja Kaligariĉ, and Igor Źibera. 2014. "Geographically weighted regression of the urban heat island of a small city." *Applied Geography* no. 53:341-353. doi: <http://dx.doi.org/10.1016/j.apgeog.2014.07.001>.
- Jacobs-Crisioni, Chris, Piet Rietveld, and Eric Koomen. 2014. "The impact of spatial aggregation on urban development analyses." *Applied Geography* no. 47:46-56.
- Keramitsoglou, Iphigenia, Chris T. Kiranoudis, Giulio Ceriola, Qihao Weng, and Umamaheshwaran Rajasekar. 2011. "Identification and analysis of urban surface temperature patterns in Greater Athens, Greece, using MODIS imagery." *Remote Sensing of Environment* no. 115 (12):3080-3090. doi: <http://dx.doi.org/10.1016/j.rse.2011.06.014>.

- Kolokotroni, Maria, Michael Davies, Ben Croxford, Saiful Bhuiyan, and Anna Mavrogianni. 2010. "A validated methodology for the prediction of heating and cooling energy demand for buildings within the Urban Heat Island: Case-study of London." *Solar Energy* no. 84 (12):2246-2255. doi: <http://dx.doi.org/10.1016/j.solener.2010.08.002>.
- Lac, C., R. P. Donnelly, V. Masson, S. Pal, S. Riette, S. Donier, S. Queguiner, G. Tanguy, L. Ammoura, and I. Xueref-Remy. 2013. "CO2 dispersion modelling over Paris region within the CO2-MEGAPARIS project." *Atmos. Chem. Phys.* no. 13 (9):4941-4961. doi: 10.5194/acp-13-4941-2013.
- Lambert-Habib, M. L., J. Hidalgo, C. Fedele, A. Lemonsu, and C. Bernard. 2013. "How is climatic adaptation taken into account by legal tools? Introduction of water and vegetation by French town planning documents." *Urban Climate* no. 4:16-34. doi: <http://dx.doi.org/10.1016/j.uclim.2013.04.004>.
- LPDAAC, Land Processes Distributed Active Archive Centre. *MODIS Reprojection Tool* 2014. Available from https://lpdaac.usgs.gov/tools/modis_reprojection_tool.
- LPDAAC, Land Processes Distributed Active Archive Centre. *MODIS Products Table - MYD11A1* 2015a. Available from https://lpdaac.usgs.gov/dataset_discovery/modis/modis_products_table/myd11a1.
- LPDAAC, Land Processes Distributed Active Archive Centre. *MODIS Products Table - MYD13Q1* 2015b. Available from https://lpdaac.usgs.gov/dataset_discovery/modis/modis_products_table/myd13q1.
- Muller, C. L., L. Chapman, S. Johnston, C. Kidd, S. Illingworth, G. Foody, A. Overeem, and R. R. Leigh. 2015. "Crowdsourcing for climate and atmospheric sciences: current status and future potential." *International Journal of Climatology* no. 35 (11):3185-3203. doi: 10.1002/joc.4210.
- Muller, Catherine L., Lee Chapman, C. S. B. Grimmond, Duick T. Young, and Xiaoming Cai. 2013. "Sensors and the city: a review of urban meteorological networks." *International Journal of Climatology* no. 33 (7):1585-1600. doi: 10.1002/joc.3678.
- Oke, TR. 1987. "Boundary layer climates. 2nd." *Methuen*, 289p.
- ONS, Office for National Statistics. *Census Geography - Super Output Area (SOA)* 2011a. Available from <http://www.ons.gov.uk/ons/index.html>.
- ONS, Office for National Statistics. *Small Area Model-Based Income Estimates, 2007/08* 2011b. Available from <http://www.ons.gov.uk/ons/index.html>.
- Pereira, Iraci Miranda, and Eleonora Sad de Assis. 2013. "Urban energy consumption mapping for energy management." *Energy Policy* no. 59:257-269. doi: <http://dx.doi.org/10.1016/j.enpol.2013.03.024>.
- Pérez-Lombard, Luis, José Ortiz, and Christine Pout. 2008. "A review on buildings energy consumption information." *Energy and Buildings* no. 40 (3):394-398. doi: <http://dx.doi.org/10.1016/j.enbuild.2007.03.007>.
- Roth, M., T. R. Oke, and W. J. Emery. 1989. "Satellite-derived urban heat islands from three coastal cities and the utilization of such data in urban climatology." *International Journal of Remote Sensing* no. 10 (11):1699-1720. doi: 10.1080/01431168908904002.
- Santamouris, M., K. Kapsis, D. Korres, I. Livada, C. Pavlou, and M. N. Assimakopoulos. 2007. "On the relation between the energy and social characteristics of the residential sector." *Energy and Buildings* no. 39 (8):893-905. doi: <http://dx.doi.org/10.1016/j.enbuild.2006.11.001>.
- Santamouris, M., N. Papanikolaou, I. Livada, I. Koronakis, C. Georgakis, A. Argiriou, and D. N. Assimakopoulos. 2001. "On the impact of urban climate on the energy consumption of buildings." *Solar Energy* no. 70 (3):201-216. doi: [http://dx.doi.org/10.1016/S0038-092X\(00\)00095-5](http://dx.doi.org/10.1016/S0038-092X(00)00095-5).

- Schwarz, Nina, Sven Lautenbach, and Ralf Seppelt. 2011. "Exploring indicators for quantifying surface urban heat islands of European cities with MODIS land surface temperatures." *Remote Sensing of Environment* no. 115 (12):3175-3186. doi: <http://dx.doi.org/10.1016/j.rse.2011.07.003>.
- Skelhorn, Cynthia, Sarah Lindley, and Geoff Levermore. 2014. "The impact of vegetation types on air and surface temperatures in a temperate city: A fine scale assessment in Manchester, UK." *Landscape and Urban Planning* no. 121:129-140. doi: <http://dx.doi.org/10.1016/j.landurbplan.2013.09.012>.
- Smith, ClaireL, A. Webb, G. J. Levermore, S. J. Lindley, and K. Beswick. 2011. "Fine-scale spatial temperature patterns across a UK conurbation." *Climatic Change* no. 109 (3-4):269-286. doi: 10.1007/s10584-011-0021-0.
- Spinsanti, Laura, and Frank Ostermann. 2013. "Automated geographic context analysis for volunteered information." *Applied Geography* no. 43:36-44. doi: <http://dx.doi.org/10.1016/j.apgeog.2013.05.005>.
- Taha, Haider, Hashem Akbari, Arthur Rosenfeld, and Joe Huang. 1988. "Residential cooling loads and the urban heat island—the effects of albedo." *Building and Environment* no. 23 (4):271-283. doi: [http://dx.doi.org/10.1016/0360-1323\(88\)90033-9](http://dx.doi.org/10.1016/0360-1323(88)90033-9).
- Tomlinson, C. J., L. Chapman, J. E. Thornes, and C. J. Baker. 2012. "Derivation of Birmingham's summer surface urban heat island from MODIS satellite images." *International Journal of Climatology* no. 32 (2):214-224. doi: 10.1002/joc.2261.
- Tomlinson, C. J., T. Prieto-Lopez, R. Bassett, L. Chapman, X. M. Cai, J. E. Thornes, and C. J. Baker. 2013. "Showcasing urban heat island work in Birmingham – measuring, monitoring, modelling and more." *Weather* no. 68 (2):44-49. doi: 10.1002/wea.1998.
- Tomlinson, Charlie, Lee Chapman, John Thornes, and Christopher Baker. 2011. "Including the urban heat island in spatial heat health risk assessment strategies: a case study for Birmingham, UK." *International Journal of Health Geographics* no. 10 (1):42.
- Tomlinson, Charlie J., Lee Chapman, John E. Thornes, Christopher J. Baker, and Tatiana Prieto-Lopez. 2012. "Comparing night-time satellite land surface temperature from MODIS and ground measured air temperature across a conurbation." *Remote Sensing Letters* no. 3 (8):657-666. doi: 10.1080/01431161.2012.659354.
- USGS, United States Geological Survey. *Band designation Landsat satellites* 2010. Available from http://landsat.usgs.gov/band_designations Landsat_satellites.php.
- USGS, United States Geological Survey. *MYD11A1 (V5) – MODIS/Aqua Land Surface Temperature and Emissivity Daily L3 1km Grid SIN* 2013. Available from <http://glovis.usgs.gov/>.
- Voogt, J. A., and T. R. Oke. 2003. "Thermal remote sensing of urban climates." *Remote Sensing of Environment* no. 86 (3):370-384. doi: [http://dx.doi.org/10.1016/S0034-4257\(03\)00079-8](http://dx.doi.org/10.1016/S0034-4257(03)00079-8).
- Weng, Qihao, Dengsheng Lu, and Jacquelyn Schubring. 2004. "Estimation of land surface temperature–vegetation abundance relationship for urban heat island studies." *Remote Sensing of Environment* no. 89 (4):467-483. doi: <http://dx.doi.org/10.1016/j.rse.2003.11.005>.
- Wilby, Robert L. 2003. "Past and projected trends in London's urban heat island." *Weather* no. 58 (7):251-260. doi: 10.1256/wea.183.02.
- Yuan, Fei, and Marvin E. Bauer. 2007. "Comparison of impervious surface area and normalized difference vegetation index as indicators of surface urban heat island effects in Landsat imagery." *Remote Sensing of Environment* no. 106 (3):375-386. doi: <http://dx.doi.org/10.1016/j.rse.2006.09.003>.

Figure 1. Electricity consumption in the UK by sector. Others: Public administration, transport, agricultural and commercial sectors.

Figure 2. a) Landuse classes in Birmingham, b) Map of the Birmingham urban area and locations.

Figure 3. a) Averaged daytime LST, b) Averaged night-time LST, c) Averaged NDVI for Birmingham in 2006.

Figure 4. Aggregated data to MSOA for a) UHI (day), b) UHI (night), c) NDVI, d) Income, e) Residential Electricity Consumption and f) Normalized Residential Electricity Consumption

Figure 5. Scatter plot diagrams with slope, intercept and R^2 a) Income by Consumption, b) Night-time LST by Consumption, c) Daytime LST by Consumption, d) NDVI by Consumption, e) Night-time LST by Income, f) Daytime LST by Income, g) NDVI by Income, h) Night-time LST by NDVI, i) Daytime LST by NDVI, j) Night-time LST by Income Normalized Consumption, k) Daytime LST by Income Normalized Consumption and l) NDVI by Income Normalized Consumption

Figure 1. Electricity consumption in the UK by sector. Others: Public administration, transport, agricultural and commercial sectors.




Figure 2. a) Land use classes in Birmingham, b) Map of the Birmingham urban area and locations.




Figure 3. a) Averaged daytime LST, b) Averaged night-time LST, c) Averaged NDVI for Birmingham in 2006.




Figure 4. Aggregated data to MSA for a) UHI (day), b) UHI (night), c) NDVI, d) Income, e) Residential Electricity Consumption and f) Normalized Residential Electricity Consumption.















Figure 5. Scatter plot diagrams with slope, intercept and R^2 a) Income by Consumption, b) Night-time LST by Consumption , c) Daytime LST by Consumption, d) NDVI by Consumption, e) Night-time LST by Income, f) Daytime LST by Income , g) NDVI by Income , h) Night-time LST by NDVI, i) Daytime LST by NDVI, j) Night-time LST by Income Normalized Consumption, k) Daytime LST by Income Normalized Consumption and l) NDVI by Income Normalized Consumption











Table 1. Ranges of lower and middle Super Output Areas

Table 2. Correlation matrix showing Pearson Correlation Coefficients between datasets

Table 3. Correlation matrix showing Pearson Correlation Coefficients between datasets and normalized electricity consumption

Table 4. Average electricity consumption by household income, England 2005 to 2011 (kWh)

Table 1. Ranges of lower and middle Super Output Areas

Geography	Minimum population	Maximum population	Minimum number of households	Maximum number of households
LSOA	1,000	3,000	400	1,200
MSOA	5,000	15,000	2,000	6,000

Table 2. Correlation matrix showing Pearson Correlation Coefficients between datasets

	Income	Ordinary Electricity/Household	MODIS night-time LST	MODIS daytime LST	MODIS NDVI
Income	-	0.62	-0.55	-0.66	0.61
Ordinary Electricity/ Household	0.62	-	-0.43	-0.47	0.39
MODIS night-time LST	-0.55	-0.43	-	0.92	-0.69
MODIS daytime LST	-0.66	-0.47	0.92	-	-0.78
MODIS NDVI	0.61	0.39	-0.69	-0.78	-

Table 3. Correlation matrix showing Pearson Correlation Coefficients between datasets and normalized electricity consumption

P	Ordinary Electricity/Household - Normalized by Income	MODIS night- time LST	MODIS daytime LST	MODIS NDVI
Ordinary Electricity/Household - Normalized by Income	-	0.41	0.52	-0.53
MODIS night-time LST	0.41	-	0.92	-0.69
MODIS daytime LST	0.52	0.92	-	-0.78
MODIS NDVI	-0.53	-0.69	-0.78	-

Table 4. Average electricity consumption by household income, England 2005 to 2011 (kWh)

		£15,000 <£15,000	£20,000	£30,000	£40,000	£50,000	£60,000	£70,000	£100,000	£150,000 and over	Unknown	Total
		£19,999	£29,999	£39,999	£49,999	£59,999	£69,999	£99,999	£149,999			
2005	3,900	4,200	4,400	4,700	4,900	5,100	5,200	5,500	6,200	6,700	5,000	4,600
2006	3,800	4,000	4,200	4,600	4,800	5,000	5,200	5,400	6,100	6,800	4,900	4,500
2007	3,700	4,000	4,200	4,500	4,800	4,900	5,100	5,400	6,100	6,700	4,900	4,500
2008	3,500	3,800	4,000	4,300	4,600	4,700	4,900	5,200	5,900	6,600	4,700	4,300
2009	3,500	3,700	3,900	4,300	4,500	4,700	4,900	5,200	5,900	6,600	4,700	4,200
2010	3,500	3,700	3,900	4,300	4,500	4,700	4,900	5,200	6,000	6,700	4,700	4,200
2011	3,400	3,700	3,800	4,200	4,500	4,700	4,900	5,200	5,900	6,700	4,600	4,200

